

IDENTIFICATION THE RATE OF ANEMIA DURING PREGNANCY AT AYUB MEDICAL COMPLEX ABBOTTABAD

AYESHA IMTIAZ MALIK^{1*}, MOMINA HAQ², MOHSINA HAQ³, BUSHRA REHMAN⁴,
FAZLI HAMID⁵, FAYAZ AHMAD MEMON⁶, TNVEER TARA⁷, HAMID ALI KHAN⁸,
SUDHAIR ABBAS BANGASH⁹ and KHALID KHAN¹⁰

^{1*}Associate Professor Histopathology Niazi Medical and Dental College Sargodha. Corresponding Author

²Associate Professor Department of Physiology Peshawar Medical College, Ripha International University Islamabad.

³Associate Professor Department of Microbiology Peshawar Medical College, Ripha International University Islamabad.

⁴Center for Biotechnology and Microbiology University of Peshawar Kp Pakistan.

⁵Department of Biochemistry Hazara University Mansehra Pakistan.

⁶Professor Department of Medicine Muhammed Medical College and Hospital, Mirpur Khas Sindh Pakistan.

⁷Department of Medical Lab Technology National Skill University Islamabad Pakistan.

⁸Bacha Khan University Charsadda kp Pakistan.

⁹Faculty of Life Sciences, Department of Pharmacy Sarhad University of science and information technology Peshawar Kp Pakistan.

¹⁰Division of Reproduction and Genetics, Division of Life Sciences and Medicine, University of Science and Technology Anhui Hafei 230027 China.

Abstract

Anemia during pregnancy is a worldwide health problem affects around 500 million women during pregnancy .Prevalence of anemia range from 5.4% to 80% in developed and developing countries respectively. Adverse effects of maternal anemia are well known and well documented on fetus as it potentiates risk of preterm labor, low birth weight babies and neonatal mortality. It also puts mothers at risk increasing maternal mortality and morbidity, antepartum and postpartum. It was a cross-sectional prospective study which was conducted at Gynecology and Obstetrics Department of Ayub Medical Complex Abbottabad. Sampling technique was Non probability convenient sampling. Duration of study was 6 month.300 Pregnant females aged 30-45 years and with parity up to 5 were enrolled in study after taking informed consent. Anemia was classified as macrocytic, microcytic and normocytic based on MCV. Fetal wellbeing was evaluated by serial abdominal ultrasound. WHO Grading for anemia was used to assess the severity of anemia. In this study 258(86%) women were multiparous 42 were primigravida. 79% were in third trimester, 15% during second trimester and 6% in first trimester. Out of 300 patients 47% had mild (11gm %), 43% had moderate (7-9gm %) and 10% were having severe anemia with Hb <7gm%. 78% had microcytic hypochromic anemia, 12% had dimorphic pictures 12% had low RBC indices with increased red cell count so these were referred for HB electrophoresis to be screened for thalassemia trait. 58 % had monthly income of 2000-4000.78% had poor diet 22% were taking normal diet. 28% had used for preparation of iron and folate for variable period of time 1-4 months.72% never used hematinic. History of blood transfusion during pregnancy and Labor was present in 16%. From the Current Research we conclude that prevalence of anemia during pregnancy is high in our society. It can have significant impact on maternal and fetal outcome it is preventable and can be treated easily. Poverty and lack of education are the most important causes of anemia during pregnancy.

Introduction:

Anemia a condition in which hemoglobin (Hb) concentration and/or red blood cell (RBC) numbers are lower than normal and insufficient to meet an individual's physiological needs (1). Anemia is associated with increased morbidity and mortality in women and children, poor birth outcomes, decreased work productivity in adults, and impaired cognitive and behavioral development in children. Preschool children (PSC) and women of reproductive age (WRA) are particularly affected (2). Anemia during pregnancy is a worldwide health problem affects around 500 million women during pregnancy (1). Anemia affects a third of the world's population and contributes to increased morbidity and mortality, decreased work productivity, and impaired neurological development (3). Prevalence of anemia range from 5.4% to 80% in developed and developing countries respectively (4) mostly it is physiological as there is more fluid retention in body leading to dilutional anemia but the true anemia that have potential to effect pregnancy adversely are nutritional anemia of iron, B12 and folate deficiency. it does not only effects maternal mortality morbidity but also have adverse effects on fetal wellbeing that can further add to fetal mortality and morbidity (5), Hemoglobin level less than 10.5g/dl at any time of gestation is regarded as anemia (6). Adverse effects of maternal anemia are well known and well documented on fetus as it potentiates risk of preterm labor, low birth weight babies and neonatal mortality (7). It also puts mothers at risk increasing maternal mortality and morbidity antepartum and postpartum (8).

MATERIAL AND METHODS

It was a cross-sectional prospective study which was conducted at Gynaecology and Obstetrics Department of Ayub Medical Complex Abbottabad. Sampling technique was Non probability convenient sampling. Duration of study was 6 month. 300 Pregnant females aged 30-45 years and with parity up to 5 were enrolled in study after informed consent. Detailed History of bleeding, parasitic infestation, pattern of taking NSAIDS, Blood transfusion and dietary customs was taken. Their socio-economic status was assessed by asking questions related to their monthly income and expenses. Blood samples for blood counts and hemoglobin estimation were collected. Cut-off value for labeling patient as anemic was 11 gram per deciliter. Anemia was classified as macrocytic, microcytic and normocytic based on MCV .MCV< 76 taken as microcytic, MCV>98 FL as macrocytic and MCV ranges 78-98FL as Normocytic. Fetal wellbeing was evaluated by serial abdominal ultrasound. Who grading for anemia was used to assess the severity of anemia.

RESULTS:

258(86%) women were multiparous 42 were primigravida. 79% were in third trimester 15% during second trimester and 6% in first trimester. Out of 300 patients 47% had mild (11gm %), 43% had moderate (7-9gm %) and 10% were having severe anemia with Hb <7gm%. 78% had microcytic hypochromic anemia, 12% had dimorphic pictures 12% had low RBC indices with increased red cell count so these were referred for HB

electrophoresis to be screened for thalassemia trait. 58 % had monthly income of 2000-4000.78% had poor diet 22% were taking normal diet. 28% had used for preparation of iron and folate for variable period of time 1-4 months.72% never used hematinic. History of blood transfusion during pregnancy and Labor was present in 16%.

Discussion:

Study was conducted to see the prevalence of anemia in pregnant women as it is a state which predisposes a women to develop anemia due to several reasons physiological or pathological overall prevalence of anemia as reported in literature globally is 74% while in developing countries like India it is reported as 19 to 61% in Tanzania it is reported as 86%, in Nigeria 47% in Philippines again 48% and Bangladesh 47%(9,10). In our study the overall prevalence of anemia is 82% which is higher than many developing countries .if compare with the local studies a study conducted in Faisalabad in 2015 reported the prevalence of anemia in pregnant woman as 75% (10,11). And study published from Lahore in 2007 reported it as 80 %.(12, 13). So our results are comparable to both of the local studies and slightly higher prevalence of anemia as most of the patients in our study were from lower socioeconomic background. 38% women had mild anemia with Hb Ranging from 8-9.9gm/dl. Out of 300 patients 47% had mild, 43% had moderate and 10% were having severe anemia. These results are comparable to a study conducted on 400 subjects from selected hospitals of Punjab which reported 44.75% patients falling in mild category, 44.50% in moderate category and 4.75% in severe category of anemia. (12). As compared to this study we have higher percentage of patients falling in severe category of anemia. (13). Another study conducted on 250 patients in southern Punjab 55.2% were anemic and out of these 60.14 % were moderately anemic while 39.86 % had mild anemia and none of these had severe anemia. Study in Faisalabad reported the mildly anemic women 24%, moderately anemic 33% and severely anemic 18% (14, 15). So in this study the percentage of severely anemic pregnant women was higher than our findings.

Among the patients which were anemic 78% had microcytic hypochromic anemia, 12% had dimorphic pictures, 12% had low MCV MCH but normal MCHC and these were referred for HB electrophoresis. The presence of anemia along with microcytic hypochromic indices indicates iron deficiency which is highest as a cause of anemia in developing countries these results are in line with the other studies .Iron deficiency anemia is already worldwide commonest cause of anemia in pregnancy. Estimated prevalence of IDA from data of 107 countries was 43% two decade back and 38% in 2011 which is much higher in developing countries 56.4%.13 We referred 12% patients for Hb Electrophoresis for Thalassemia trait as it is also important to diagnose it to reduce the burden of this disease as our part of the world is already affected by it. 12% patients had dimorphic population of RBCs these patients were later found to have mixed nutritional deficiency which is very much near to the expectations as most of the patients in our study were from lower socioeconomic group having very limited incomes also their diet was not satisfactory and lacking in many important nutrients ,proteins and

vitamins. Mixed nutritional deficiency that is iron and folate both of which are required in pregnancy can worsen the condition in fetus and in mother as well. Folate deficiency during pregnancy can result in congenital birth defects (16). 72% patients in our study were totally ignorant about the use of hematinic considering them unimportant and of no significance. Hook worm infestation and nutritional deficiency were the most important factors in these anemic patients. A study conducted in India also observed the two factors as most important cause of anemia (17)16% patients had transfusion history repeatedly during their pregnancy for severe anemia.

Conclusion:

We consider that education and awareness is very important for prevention and treatment of anemia as it has worse outcome for maternal and fetal health. Poverty and ignorance are two very important reason for anemia and poor health of woman during pregnancy. Worthy side is that it is not only treatable but also preventable by simply paying attention to the normal dietary patterns and general wellbeing of the mother. Regular use of hematinic which are routinely prescribed in pregnancy can also prevent the unwanted and undesirable outcomes.

References:

- Young MF. Maternal anaemia and risk of mortality: a call for action. *The Lancet Global Health*. 2018 May 1;6(5):e479-80
- Bora R, Sable C, Wolfson J, et al. Prevalence of anemia in pregnant women and its effect on neonatal outcomes in Northeast India. *J Matern Fetal Neonatal Med*. 2014;27:887–891
- Noronha JA, Al Khasawneh E, Seshan V, Ramasubramaniam S, Raman S. Anemia in pregnancy-consequences and challenges: a review of literature. *Journal of South Asian Federation of Obstetrics and Gynecology*. 2012 Jan;4(1):64-70
- Breymann C. Iron deficiency anemia in pregnancy. In *Seminars in hematology* 2015 Oct 1 (Vol. 52, No. 4, pp. 339-347). WB Saunders.
- Rahman MM, Abe SK, Rahman MS, Kanda M, Narita S, Bilano V, Ota E, Gilmour S, Shibuya K. Maternal anemia and risk of adverse birth and health outcomes in low-and middle-income countries: systematic review and meta-analysis, 2. *The American journal of clinical nutrition*. 2016 Feb 1; 103(2):495-504.
- Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, De Onis M, Ezzati M, Grantham-McGregor S, Katz J, Martorell R, Uauy R. Maternal and child undernutrition and overweight in low-income and middle-income countries. *The lancet*. 2013 Aug 3;382(9890):427-51.
- Prasad KN. Prevalence of anemia among pregnant women-a cross-sectional study. *International Journal of Medical Science and Public Health*. 2018;7(12):1023-6
- Anjum A, Manzoor M, Manzoor N, Shakir HA. Prevalence of anemia during pregnancy in district Faisalabad, Pakistan. *Punjab Univ J Zool*. 2015;30(1):15-20.
- KHAN, Z.M., SHOAIB, M. AND KHALID, M., 2007. Anemia during antenatal period; evaluation of different related parameters in pregnant women. *Professional Med. J. Mar.*,14(1): 1-6.

Garg SK, Kaur H, Kaur H, Sharma J, Kaur J, Kaur K, Kaur K, Kaur M. Prevalence of Anemia among Antenatal mothers attending antenatal OPD's in selected hospitals of Punjab. International Journal of Nursing Education and Research. 2020;8(2):175-8.

SAFDAR S, Mirbahar A, AWAN Z. Anemia in pregnancy: Related risk factors in under developed area. The Professional Medical Journal. 2011 Mar 10;18(01):1-4.

Anjum A, Manzoor M, Manzoor N, Shakir HA. Prevalence of anemia during pregnancy in district Faisalabad, Pakistan. Punjab Univ J Zool. 2015;30(1):15-20.

Tandon R, Jain A, Malhotra P. Management of iron deficiency anemia in pregnancy in India. Indian Journal of Hematology and Blood Transfusion. 2018 Apr;34(2):204-15.

Bhide P. Neural Tube Defects and Folate Status in India. In Birth Defects in India 2021 (pp. 235-249). Springer, Singapore

Vasanthamani P, Thiruvankideswary B, Meena TS, Padmanaban S. Maternal and fetal outcome in anaemia complicating pregnancy.